

# MA 225 Test 1

SEPTEMBER 22, 2010

Dr. Franke

Name \_\_\_\_\_

Show all work. You may not use a calculator.

- (15%) Given the statement: No isosceles triangle is an equilateral triangle. (Universe is all triangles.)
  - translate the statement into symbols with quantifiers.
  - write in symbols a useful denial of your statement in (a.)
  - translate your answer in (b.) back into English.
- (5%) Define what is meant by a counterexample to  $(\forall x)P(x) \Rightarrow Q(x)$ .
- (8%) Outline a proof of  $P \Rightarrow Q \vee R$  that uses contraposition.
- (10%) Given the proposition: If  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is an  $x \in (a, b)$  so that

$$f'(x) = \frac{f(b) - f(a)}{b - a}.$$

- Write the converse.
  - Write the contrapositive.
- (8%) Find the truth table for  $(P \vee \sim Q) \Rightarrow \sim (P \wedge Q)$ . Determine if this proposition is a tautology, a contradiction or neither. (explain)
  - (24%) For each of the following, determine if it is true or false.
    - $(\exists x)(\forall y)(2x = y)$  (assume the universe is the real numbers)
    - $(\forall y)(\exists x)(x = \sin(y))$  (assume the universe is the real numbers)
    - $(\forall x)(\exists y)(xy \text{ is odd})$  (assume the universe is the integers)
    - $(\forall y)(\exists! x)(y = x^3)$  (assume the universe is the real numbers)
    - If  $P(x)$  is an open statement then  $\sim [(\exists! x)P(x)]$  is equivalent to  $[(\forall x)(\sim P(x))] \vee [(\exists y)(\exists z)[P(y) \wedge P(z) \wedge (y \neq z)]]$ .
    - $P \wedge Q \Rightarrow R$  is equivalent to  $(P \Rightarrow R) \wedge (Q \Rightarrow R)$ .
  - (30%) Prove each of the following:
    - If  $n$  is an integer, then  $5n^2 + 3n$  is an even number. (Use 2 cases.)
    - If  $a, b, c$  and  $d$  are natural numbers,  $a$  divides  $c$  and  $b$  divides  $d$ , then  $ab$  divides  $cd$ .

1.  $\frac{5}{5} (\forall x)$   $x$  is isosceles  $\Rightarrow x$  is not equilateral

$\frac{5}{5} (\exists x)$   $x$  is isosceles and  $x$  is equilateral

$\frac{5}{5}$  Some isosceles triangle is an equilateral triangle.

2.  $\frac{5}{5}$  A counterexample is an  $x$  where  $P(x)$  and  $\sim Q(x)$  are both true

3. Assume  $\sim Q$  and  $\sim R$

$\frac{5}{5}$   $\vdots$

Thus  $\sim P$ .

Hence  $\sim Q \wedge \sim R \Rightarrow \sim P$

Thus by contraposition  $P \Rightarrow Q \vee R$ .


4. a.)  $\frac{5}{5}$  If there is an  $x \in (a, b)$  so that  $f'(x) = \frac{f(b) - f(a)}{b - a}$

then  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ .

b.)  $\frac{5}{5}$  If  $\forall x \in (a, b)$   $f'(x) \neq \frac{f(b) - f(a)}{b - a}$  then  $f$  is not continuous on  $[a, b]$  or  $f$  is not differentiable on  $(a, b)$ .

5.

P	Q	$\sim Q$	$P \wedge Q$	$\sim(P \wedge Q)$	$P \vee \sim Q$	$(P \vee \sim Q) \leftrightarrow \sim(P \wedge Q)$
T	T	F	T	F	T	F
F	T	F	F	T	F	T
T	F	T	F	T	T	T
F	F	T	F	T	T	T



not all T so not a tautology

not all F so not a contradiction

6

- a F
- b T
- c F
- d T
- e T
- f F

7a.) Let  $n$  be an integer. Either  $n$  is even or  $n$  is odd.

[assume  $n$  is even. Then  $\exists k \in \mathbb{Z} \ni n = 2k$ ,

$$5n^2 + 3n = 5(2k)^2 + 3(2k) = 2(10k^2 + 3k).$$

$10k^2 + 3k \in \mathbb{Z}$  by closure. Thus  $5n^2 + 3n$  is even.

Thus if  $n$  is even then  $5n^2 + 3n$  is even.

[assume  $n$  is odd. Then  $\exists l \in \mathbb{Z} \ni n = 2l + 1$ ,

$$5n^2 + 3n = 5(2l+1)^2 + 3(2l+1) = 5(4l^2 + 4l + 1) + 3(2l+1)$$

$$= 20l^2 + 20l + 5 + 6l + 3 = 20l^2 + 26l + 8$$

$$= 2(10l^2 + 13l + 4). \quad \underline{10l^2 + 13l + 4 \in \mathbb{Z} \text{ by closure.}}$$

Thus  $5n^2 + 3n$  is even.

Hence if  $n$  is odd,  $5n^2 + 3n$  is even.

[since we have checked both cases  $n$  even and  $n$  odd,  
if  $n$  is an integer then  $5n^2 + 3n$  is even.]

7. b. Let  $a, b, c, d$  be natural numbers. Assume  $a$  divides  $c$   
and  $b$  divides  $d$ . Then  $\exists k \in \mathbb{N}$  and  $l \in \mathbb{N}$   $\Rightarrow$

$$\underline{c = ak}, \text{ and } \underline{d = bl}. \text{ Thus}$$

$$cd = (ak)(bl) = ab(kl). \quad \underline{kl \in \mathbb{N} \text{ by closure.}}$$

Thus  $ab$  divides  $cd$ .

[Therefore if  $a, b, c, d$  are natural numbers,  $a$  divides  $c$   
and  $b$  divides  $d$ , then  $ab$  divides  $cd$ .