

# Mahonian pairs

Bruce E. Sagan

Department of Mathematics, Michigan State University,  
East Lansing, MI 48824-1027, USA, [sagan@math.msu.edu](mailto:sagan@math.msu.edu)

and

Carla D. Savage

Department of Computer Science, North Carolina State University,  
Raleigh, NC 27695-8206, USA, [savage@ncsu.edu](mailto:savage@ncsu.edu)

January 22, 2011

Key Words: ballot sequence, Greene-Kleitman symmetric chain decomposition, Foata's fundamental bijection, integer partition, inversion number, Mahonian statistic, major index, rank of a partition,  $q$ -Catalan number,  $q$ -Fibonacci number

AMS subject classification (2000): Primary 05A05; Secondary 05A10, 05A15, 05A19, 05A30, 11P81.

## Abstract

We introduce the notion of a Mahonian pair. Consider the set,  $\mathbb{P}^*$ , of all words having the positive integers as alphabet. Given finite subsets  $S, T \subset \mathbb{P}^*$ , we say that  $(S, T)$  is a *Mahonian pair* if the distribution of the major index,  $\text{maj}$ , over  $S$  is the same as the distribution of the inversion number,  $\text{inv}$ , over  $T$ . So the well-known fact that  $\text{maj}$  and  $\text{inv}$  are equidistributed over the symmetric group,  $\mathfrak{S}_n$ , can be expressed by saying that  $(\mathfrak{S}_n, \mathfrak{S}_n)$  is a Mahonian pair. We investigate various Mahonian pairs  $(S, T)$  with  $S \neq T$ . Our principal tool is Foata's fundamental bijection  $\phi : \mathbb{P}^* \rightarrow \mathbb{P}^*$  since it has the property that  $\text{maj } w = \text{inv } \phi(w)$  for any word  $w$ . We consider various families of words associated with Catalan and Fibonacci numbers. We show that, when restricted to words in  $\{1, 2\}^*$ ,  $\phi$  transforms familiar statistics on words into natural statistics on integer partitions such as the size of the Durfee square. The Rogers-Ramanujan identities, the Catalan triangle, and various  $q$ -analogues also make an appearance. We generalize the definition of Mahonian pairs to infinite sets and use this as a tool to connect a partition bijection of Corteel-Savage-Venkatraman with the Greene-Kleitman decomposition of a Boolean algebra into symmetric chains. We close with comments about future work and open problems.

# 1 Introduction

To introduce our principal object of study, Mahonian pairs, we need to set up some notation. Let  $\mathbb{N}$  and  $\mathbb{P}$  denote the nonnegative and positive integers, respectively. Consider the Kleene closure  $\mathbb{P}^*$  of all words  $w = a_1 a_2 \dots a_n$  where  $a_i \in \mathbb{P}$  for  $1 \leq i \leq n$  and  $n \geq 0$ . We let  $|w| = n$  be the length of  $w$  and  $\epsilon$  be the empty word of length 0. We will often express words using multiplicity notation where  $w = a^{m_1} b^{m_2} \dots c^{m_k}$  is the word beginning with  $m_1$  copies of  $a$  followed by  $m_2$  copies of  $b$ , and so forth. Denote by  $\Pi(w)$  the subset of  $\mathbb{P}^*$  consisting of all permutations of  $w$ .

We will be considering various well-known statistics on  $\mathbb{P}^*$ . The word  $w = a_1 a_2 \dots a_n$  has *descent set*

$$\text{Des } w = \{i : 1 \leq i < n \text{ and } a_i > a_{i+1}\},$$

and *inversion set*

$$\text{Inv } w = \{(i, j) : 1 \leq i < j \leq n \text{ and } a_i > a_j\},$$

and we say that  $a_i$  is *in inversion with*  $a_j$  if  $(i, j) \in \text{Inv } w$ . From these sets we get the *major index*

$$\text{maj } w = \sum_{i \in \text{Des } w} i,$$

the *descent number*

$$\text{des } w = \# \text{Des } w,$$

and the *inversion number*

$$\text{inv } w = \# \text{Inv } w$$

where the pound sign indicates cardinality. For more information on these statistics, see Stanley's text [58, p. 21 & ff.].

Let  $\mathfrak{S}_n \subset \mathbb{P}^*$  be the symmetric group of all permutations of  $\{1, \dots, n\}$ . A celebrated result of MacMahon [50, pp. 508–549 and pp. 556–563] states that the distribution of  $\text{maj}$  and  $\text{inv}$  over  $\mathfrak{S}_n$  are the same, i.e., we have equality of the generating functions

$$\sum_{w \in \mathfrak{S}_n} q^{\text{maj } w} = \sum_{w \in \mathfrak{S}_n} q^{\text{inv } w}. \quad (1)$$

A statistic  $\text{st} : \mathfrak{S}_n \rightarrow \mathbb{N}$  is called *Mahonian* if it is equidistributed with  $\text{maj}$  and  $\text{inv}$  and there is a large literature on such statistics. This motivates our new definition. Given finite subsets  $S, T \subseteq \mathbb{P}^*$ , we call  $(S, T)$  a *Mahonian pair* if

$$\sum_{w \in S} q^{\text{maj } w} = \sum_{w \in T} q^{\text{inv } w} \quad (2)$$

So equation (1) can be expressed by saying that  $(\mathfrak{S}_n, \mathfrak{S}_n)$  is a Mahonian pair.

Note that the Mahonian pair relation is not symmetric. We wish to study various pairs where  $S \neq T$ . Our main tool will be Foata's fundamental bijection [33]  $\phi : \mathbb{P}^* \rightarrow \mathbb{P}^*$  (defined in the next section) because it has the property that

$$\text{maj } w = \text{inv } \phi(w) \tag{3}$$

for all  $w \in \mathbb{P}^*$ . So for any finite  $S \subseteq \mathbb{P}^*$  we have corresponding Mahonian pairs  $(S, \phi(S))$  and  $(\phi^{-1}(S), S)$ . Of course, the point of the definition (2) is that  $S$  and  $T$  should have independent interest outside of being part of a pair.

The rest of this paper is structured as follows. In the next section we recall the definition of Foata's bijection,  $\phi$ , and prove some general results which will be useful in the sequel. One of the goals of this work is to show that, when restricted to  $\{1, 2\}^*$ ,  $\phi$  sends various well-known statistics on words to natural statistics on partitions; see, for example, equation (15). Section 3 studies pairs associated with ballot sequences and Catalan numbers. Pairs connected with Fibonacci numbers are studied in Section 4. The definition of Mahonian pairs is generalized to infinite sets in Section 5 and we use this idea to connect a bijection of Corteel-Savage-Venkatraman [29] with the Greene-Kleitman decomposition of a Boolean algebra into symmetric chains [40]. The final section is devoted to remarks and open problems.

## 2 Foata's fundamental bijection

We now review the construction of Foata's bijection  $\phi : \mathbb{P}^* \rightarrow \mathbb{P}^*$ . Given  $v = a_1 a_2 \dots a_n$  we inductively construct a sequence of words  $w_1, w_2, \dots, w_n = \phi(v)$  as follows. Let  $w_1 = a_1$ . To form  $w_{i+1}$  from  $w_i = b_1 b_2 \dots b_i$ , compare  $a_{i+1}$  with  $b_i$ . Form the unique factorization  $w_i = f_1 f_2 \dots f_k$  such that, if  $b_i \leq a_{i+1}$  (respectively,  $b_i > a_{i+1}$ ), then each factor contains only elements greater than (respectively, less than or equal to)  $a_{i+1}$  except the last which is less than or equal to (respectively, greater than)  $a_{i+1}$ . Let  $g_j$  be the cyclic shift of  $f_j$  which brings the last element of the factor to the front and let  $w_{i+1} = g_1 g_2 \dots g_k a_{i+1}$ . As an example, finding  $\phi(2121312)$  would give rise to the following computation where we separate the

factors of  $w_i$  with dots:

$$\begin{aligned}
w_1 &= 2 &= 2 && \text{since } a_2 = 1, \\
w_2 &= 21 &= 2 \cdot 1 && \text{since } a_3 = 2, \\
w_3 &= 212 &= 2 \cdot 12 && \text{since } a_4 = 1, \\
w_4 &= 2211 &= 2 \cdot 2 \cdot 1 \cdot 1 && \text{since } a_5 = 3, \\
w_5 &= 22113 &= 2 \cdot 2 \cdot 113 && \text{since } a_6 = 1, \\
w_6 &= 223111 &= 2 \cdot 2 \cdot 31 \cdot 1 \cdot 1 && \text{since } a_7 = 2, \\
w_7 &= 2213112 &= \phi(2121312).
\end{aligned}$$

One can show that  $\phi$  is a bijection by constructing a step-by-step inverse. Equation (3) can be shown to hold by noting that  $w_i = \phi(v_i)$  where  $v_i = a_1 a_2 \dots a_i$  and then showing that at each stage of the algorithm the change in maj in passing from  $v_i$  to  $v_{i+1}$  is the same as the change in inv in passing from  $w_i$  to  $w_{i+1}$ .

We will be particularly interested in the action of  $\phi$  on words  $w \in \{1, 2\}^*$ . Keeping the notation of the previous paragraph, note that if  $a_{i+1} = 2$  then all the factors are of length 1 and so  $w_{i+1} = w_i 2$ . If  $a_{i+1} = 1$  then the factors will either be of the form  $1^n 2$  if  $b_i = 2$  or of the form  $2^n 1$  if  $b_i = 1$  for some  $n \geq 0$ . This enables us to give a nice characterization of  $\phi$ . In the proof of this result and later on, it will be convenient to have another notation for words in  $\{1, 2\}^*$  by subscripting the ones right to left with  $1, 2, \dots$  and the twos similarly left to right. For example,  $w = 112122$  would become  $w = 1_3 1_2 2_1 1_1 2_2 2_3$ .

**Lemma 2.1.** *The map  $\phi$  on  $\{1, 2\}^*$  can be defined recursively by  $\phi(\epsilon) = \epsilon$ ,  $\phi(1) = 1$  and the three rules*

- (i)  $\phi(w2) = \phi(w)2$ ,
- (ii)  $\phi(w11) = 1\phi(w1)$ ,
- (iii)  $\phi(w21) = 2\phi(w)1$ .

**Proof.** Since the initial conditions and recursive rules uniquely define a map  $\phi' : \{1, 2\}^* \rightarrow \{1, 2\}^*$ , it suffices to show that  $\phi$  satisfies these statements to show that  $\phi' = \phi$ . The initial conditions and (i) follow directly from the description of  $\phi$  given above.

(ii) Let  $v = \phi(w1)$ . To form  $\phi(w11)$ , one must cycle the factors of  $v$  which are all of the form  $2^n 1$  for some  $n$ . During this process,  $1_i$  in  $v$  moves into the position of  $1_{i+1}$  for all  $i$  except when  $i$  has attained its maximum. In that case,  $1_i$  becomes a new one at the beginning of the resulting word. And the final one in  $w11$  takes the place of  $1_1$  in  $v$ . Thus the total effect is to move  $v$  over one position and prepend a one, in other words  $\phi(w11) = 1\phi(w1)$ .

(iii) The proof is similar to that of (ii) while also using (i). So it is left to the reader.  $\square$

Note that  $v \in \{1, 2\}^*$  has  $\text{des } v = d$  if and only if one can write

$$v = 1^{m_0} 2^{n_0} 1^{m_1} 2^{n_1} \dots 1^{m_d} 2^{n_d} \quad (4)$$

where  $m_0, n_d \geq 0$  and  $m_i, n_j > 0$  for all other  $i, j$ .

**Proposition 2.2.** *Let  $v$  have  $d$  descents and so be given by (4). It follows that*

$$\phi(v) = 1^{m_d-1} 2 1^{m_{d-1}-1} 2 \dots 1^{m_1-1} 2 1^{m_0} 2^{n_0-1} 1 2^{n_1-1} 1 \dots 2^{n_{d-1}-1} 1 2^{n_d}.$$

**Proof.** We induct on  $d$ . When  $d = 0$  we have  $v = 1^{m_0} 2^{m_0} = \phi(v)$  which is correct. For  $d > 1$ , one can write

$$v = u 2 1^{m_d} 2^{n_d}$$

where  $u$  is the appropriate prefix of  $v$ . Using the previous lemma repeatedly gives

$$\begin{aligned} \phi(v) &= \phi(u 2 1^{m_d} 2^{n_d}) \\ &= 1^{m_d-1} \phi(u 2 1) 2^{n_d} \\ &= 1^{m_d-1} 2 \phi(u) 1 2^{n_d}. \end{aligned}$$

By induction, this last expression coincides with the desired one.  $\square$

Applying  $\phi^{-1}$  to both sides of the last set of displayed equations, one gets the following result which we record for later use.

**Lemma 2.3.** *If  $w = 1^m 2 u 1 2^n$  for  $m, n \geq 0$  then*

$$\phi^{-1}(w) = \phi^{-1}(u) 2 1^{m+1} 2^n. \quad \square$$

There is an intimate connection between words in  $\{1, 2\}^*$  and integer partitions. An *integer partition* is a weakly decreasing sequence  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$  of positive integers. A cornucopia of information about partitions can be found in Andrews' book [4]. The  $\lambda_i$  are called *parts* and we will use multiplicity notation for them as we do in words. Let  $|\lambda|$  denote the sum of its parts. The *Ferrers diagram* of  $\lambda$  consists of  $k$  left-justified rows of boxes with  $\lambda_i$  boxes in row  $i$ . The Ferrers diagram for  $\lambda = (3, 2, 2) = (3, 2^2)$  is shown northwest of the dark path in Figure 1. We let  $(i, j)$  denote the box in row  $i$  and column  $j$ . We say that  $\lambda$  *fits in an*  $m \times n$  *rectangle*, written  $\lambda \subseteq m \times n$ , if  $\lambda \subseteq (m^n)$  as Ferrers diagrams. Figure 1 shows that  $(3, 2, 2) \subseteq 5 \times 4$ . The *Durfee square* of  $\lambda$ ,  $D(\lambda)$ , is the largest partition  $(d^d) \subseteq \lambda$ . We let  $d(\lambda) = d$  denote the length of a side of  $D(\lambda)$  and in our example




Figure 1: A partition  $\lambda$  contained in  $5 \times 4$  and the corresponding lattice path

$d(3, 2, 2) = 2$ . In general, if  $\lambda \subseteq \mu$  then we have a *skew partition*  $\lambda/\mu$  consisting of all the boxes of the Ferrers diagram of  $\lambda$  which are not in  $\mu$ . So the Ferrers diagram of  $(4^5)/(3, 2^2)$  consists of the boxes southeast of the dark path in the figure.

Consider any word  $w$  in the set  $\Pi(1^m 2^n)$  of permutations of  $1^m 2^n$ . We identify  $w$  with the a lattice path  $P(w)$  in  $\mathbb{Z}^2$  from  $(0, 0)$  to  $(n, m)$ , where a 1 denotes a step one unit north and a 2 a step one unit east. Figure 1 displays the path for  $w = 112211212$  with each step labeled by the corresponding element of  $w$  along with its subscript. In this way we associate with  $w$  an integer partition  $\lambda(w) \subseteq m \times n$  whose Ferrers diagram consists of the boxes inside the rectangle and northwest of  $P(w)$ . Note that  $2_j$  is in inversion with  $1_i$  if and only if the Ferrers diagram of  $\lambda(w)$  contains the box  $(i, j)$ . It follows that

$$\text{inv } w = |\lambda(w)|. \quad (5)$$

Also,  $d(\lambda)$  is the largest subscript such that  $2_d$  is in inversion with  $1_d$ .

The following result will be useful in deriving generating functions using  $\phi$ .

**Corollary 2.4.** *If  $v \in \{1, 2\}^*$  and  $\lambda = \lambda(\phi(v))$ , then*

- (i)  $\text{maj } v = |\lambda|$ , and
- (ii)  $\text{des } v = d(\lambda)$ .

**Proof.** Let  $w = \phi(v)$ . By the properties of Foata's map and (5) we have  $\text{maj } v = \text{inv } w = |\lambda|$ .

For (ii), suppose  $v$  has the form (4). From Proposition 2.2 we can read off the positions of the first  $d$  ones and the last  $d$  twos in  $w = \phi(v)$ . In particular,  $2_d$  is in inversion with  $1_d$  and this is not true for any larger subscript. Thus  $d(\lambda) = d = \text{des } v$  as desired.  $\square$

### 3 Ballot sequences

#### 3.1 Applying $\phi$

We will now investigate the effects of  $\phi$  and  $\phi^{-1}$  on ballot sequences. Say that  $w \in \mathbb{P}^*$  is a *ballot sequence* if, for every prefix  $v$  of  $w$  and every  $i \in \mathbb{P}$ , the number of  $i$ 's in  $w$  is at least as large as the number of  $(i + 1)$ 's. Consider

$$B_n = \{w \in \Pi(1^n 2^n) : w \text{ is a ballot sequence}\}.$$

The number of such ballot sequence is given by the *Catalan number*

$$C_n = \frac{1}{n+1} \binom{2n}{n}. \tag{6}$$

There is a corresponding  $q, t$ -analogue

$$c_n(q, t) = \sum_{w \in B_n} q^{\text{maj } w} t^{\text{des } w}. \tag{7}$$

The  $c_n(q, t)$  were first defined in a paper of F\"urlinger and Hofbauer [34]. The case  $t = 1$  was the subject of an earlier note by Aissen [1]. These  $q$ -analogues have since been studied by various authors [17, 18, 46].

In order to describe  $\phi(B_n)$ , we need to define the ranks of a partition,  $\lambda$ . We let the  $i$ th rank of  $\lambda$ ,  $1 \leq i \leq d(\lambda)$ , be

$$r_i(\lambda) = \lambda_i - \lambda'_i$$

where  $\lambda'$  is the *conjugate* of  $\lambda$  obtained by transposing  $\lambda$ 's Ferrers diagram. We also let

$$R_n = \{\lambda : \lambda \subseteq n \times n \text{ and } r_i(\lambda) < 0 \text{ for all } 1 \leq i \leq d(\lambda)\}.$$

The concept of rank goes back to Dyson [30]. Partitions with all ranks positive (which are in bijection with partitions with all ranks negative by conjugation) have arisen in the work of a number of authors [3, 6, 32, 54]. Interest in them stems from connections with graphical partitions and with the Rogers-Ramanujan identities. It follows from the work of Corteel, Savage, and Venkatraman [29] that  $\#R_n = C_n$  which is also a corollary of the next result.

**Theorem 3.1.** *We have*

$$\phi(B_n) = \{w \in \Pi(1^n 2^n) : \lambda(w) \in R_n\}, \tag{8}$$

and so

$$c_n(q, t) = \sum_{\lambda \in R_n} q^{|\lambda|} t^{d(\lambda)}.$$

**Proof.** We first prove equation (8). Assume that  $v \in B_n$  has the form (4). Then  $v \in B_n$  is equivalent to  $m_0 + \cdots + m_k \geq n_0 + \cdots + n_k$  for  $0 \leq k \leq d$  with both sides equalling  $n$  for  $k = d$ .

To check the ranks of  $\lambda(w)$ , note that from our description of the relationship between  $w$  and  $\lambda(w)$  it follows that  $\lambda_i$  is the number of twos before  $1_i$  in  $w$ , and  $\lambda'_j$  is the number of ones after  $2_j$ . Using Proposition 2.2, we have

$$\begin{aligned}\lambda_l &= d + (n_0 - 1) + (n_1 - 1) + \cdots + (n_{d-l} - 1), \\ \lambda'_l &= d + m_0 + (m_1 - 1) + \cdots + (m_{d-l} - 1)\end{aligned}$$

for  $1 \leq l \leq d$ . Hence  $r_l(\lambda(w)) < 0$  if and only if  $\lambda'_l > \lambda_l$  for  $1 \leq l \leq d$ , which is clearly equivalent to the ballot conditions for  $0 \leq k < d$ . And we have the desired equality when  $k = d$  since  $w \in \Pi(1^n 2^n)$ ,

The generating function identity follows directly from (8) and Corollary 2.4. This finishes the proof of the theorem.  $\square$

### 3.2 Applying $\phi^{-1}$

We can obtain another Mahonian pair by applying  $\phi^{-1}$  to  $B_n$ . For the proof characterizing the preimage, we need a notion of conjugation for sequences. If  $w = b_1 b_2 \dots b_n \in \{1, 2\}^*$  then let

$$w' = b'_n \dots b'_2 b'_1 \tag{9}$$

where  $b'_i = 3 - b_i$  for  $1 \leq i \leq n$ , i.e., read the sequence backwards while exchanging the ones and twos. It should be clear from the definitions that  $\lambda(w)$  and  $\lambda(w')$  are conjugate partitions.

**Theorem 3.2.** *Let  $v$  have the form (4). Then  $\phi(v) \in B_n$  if and only if we have the conditions:*

(i) for all  $i$  with  $1 \leq i \leq d$

$$\begin{aligned}m_d + m_{d-1} + \cdots + m_{d-i+1} &\geq 2i, \\ n_d + n_{d-1} + \cdots + n_{d-i+1} &\geq 2i - 1,\end{aligned}$$

(ii) and

$$\sum_{i=0}^d m_i = \sum_{i=0}^d n_i = n.$$



**Proof.** Suppose  $\phi(v) \in B_n$ . Then, using Proposition 2.2, we see that the ballot sequence condition for the prefix up to  $2_i$  (where  $1 \leq i \leq d$ ) is given by

$$(m_d - 1) + (m_{d-1} - 1) + \cdots + (m_{d-i+1} - 1) \geq i.$$

This, in turn, is clearly equivalent to the first inequality in (i).

To obtain the second inequality, note that if  $w$  is in  $B_n$  then so is  $w'$  as defined above. Now use the same reasoning as in the previous paragraph.

Finally, condition (ii) follows since  $\phi$  preserves the number of ones and twos. It is easy to see that the reasoning above is reversible and so (i) and (ii) imply  $\phi(v) \in B_n$ .  $\square$

Let us put the previous theorem in context. Consider

$$B_{m,n} = \{w \in \Pi(1^m 2^n) : w \text{ is a ballot sequence}\}$$

The *Catalan triangle* (Sloane A008315) has entries  $C_{n,d}$  for  $0 \leq d \leq \lfloor n/2 \rfloor$ , where

$$C_{n,d} = \#B_{n-d,d}.$$

It is well known that

$$C_n = \sum_{d \geq 0} C_{n,d}^2 \tag{10}$$

where we assume  $C_{n,d} = 0$  for  $d > \lfloor n/2 \rfloor$ . One way to see this is to consider the map  $\beta : B_n \rightarrow \uplus_{d \geq 0} B_{n-d,d}^2$  defined as follows. Given  $w = b_1 b_2 \dots b_{2n} \in B_n$ , write  $w = xy$  where  $x = b_1 b_2 \dots b_n$  and  $y = b_{n+1} b_{n+2} \dots b_{2n}$ , and define

$$\beta(xy) = (x, y') \tag{11}$$

where  $y'$  is as in (9). It is easy to show that  $\beta$  is well defined and a bijection and so (10) follows.

Associated with any  $v$  of the form (4) are two compositions (ordered partitions allowing zeros), the *one's composition*  $\omega(v) = (m_0, m_1, \dots, m_d)$  and the *two's composition*  $\tau(v) = (n_0, n_1, \dots, n_d)$ . Note that  $\omega(v) \in \mathbb{N}\mathbb{P}^d$  while  $\tau(v) \in \mathbb{P}^d\mathbb{N}$ . Consider two sets of compositions

$$O_{n,d} = \{\omega \in \mathbb{N}\mathbb{P}^d : |\omega| = n \text{ and } \omega_d + \omega_{d-1} + \cdots + \omega_{d-i+1} \geq 2i \text{ for } 1 \leq i \leq d\},$$

and

$$T_{n,d} = \{\tau \in \mathbb{P}^d\mathbb{N} : |\tau| = n \text{ and } \tau_d + \tau_{d-1} + \cdots + \tau_{d-i+1} \geq 2i - 1 \text{ for } 1 \leq i \leq d\}.$$

To describe the next result, let  $A_{n,d}$  be the set of  $v$  of the form (4) such that  $\phi(v) \in B_n$ . Also note that if we define a map  $f$  on  $A_{n,d}$  for each  $d \geq 0$ , then  $f$  can be considered as a map on  $\uplus_{d \geq 0} A_{n,d} = \phi^{-1}(B_n)$ . Similar considerations apply to the other sets with subscripts  $n, d$  defined above.

**Theorem 3.3.** *We have the following facts.*

(i) *The maps  $o : O_{n,d} \rightarrow B_{n-d,d}$  and  $t : T_{n,d} \rightarrow B_{n-d,d}$  defined by*

$$o(\omega_0, \omega_1, \dots, \omega_d) = 1^{\omega_d-1} 2 1^{\omega_{d-1}-1} 2 \dots 1^{\omega_1-1} 2 1^{\omega_0},$$

and

$$t(\tau_0, \tau_1, \dots, \tau_d) = 1^{\tau_d} 2 1^{\tau_{d-1}-1} 2 1^{\tau_{d-2}-1} \dots 2 1^{\tau_0-1}$$

are bijections. Thus

$$\#O_{n,d} = \#T_{n,d} = C_{n,d}.$$

(ii) *The map  $\omega \times \tau : A_{n,d} \rightarrow O_{n,d} \times T_{n,d}$  given by  $v \mapsto (\omega(v), \tau(v))$  is a bijection. Thus*

$$\#A_{n,d} = C_{n,d}^2.$$

(iii) *Composing maps from right to left, we have*

$$\beta = (o \times t) \circ (\omega \times \tau) \circ \phi^{-1}.$$

where  $\beta$  is the map given by (11).

**Proof.** (i) We will prove the statements involving  $O_{n,d}$  as proofs for  $T_{n,d}$  is similar. We must first show that  $o$  is well defined, i.e., that if  $\omega \in O_{n,d}$  then  $o(\omega) \in B_{n-d,d}$ . The number of ones in  $o(\omega)$  is  $\omega_0 + \sum_{i=1}^d (\omega_i - 1) = n - d$  as desired, and it is clear that there are exactly  $d$  twos. To verify the ballot condition, it suffices to check the prefix ending in  $2_j$  for  $1 \leq j \leq d$ . Using the defining inequality for  $O_{n,d}$ , the number of ones in this prefix is

$$\sum_{i=1}^j (\omega_{d-i+1} - 1) \geq 2j - j = j$$

which is what we need. Constructing an inverse to prove bijectivity is easy, and the statement about cardinalities follows from  $o$  being bijective.

(ii) The fact that the map is well defined and bijective is just a restatement of Theorem 3.2. The cardinality of  $A_{n,d}$  can now be computed using this bijection and (i).

(iii) Any  $w \in B_n$  can be written uniquely as  $w = xy$  where  $|x| = |y| = n$ . Say  $x \in \Pi(1^{n-d} 2^d)$  for some  $d \geq 0$ . It follows that  $y \in \Pi(1^d 2^{n-d})$ . So we can write

$$\begin{aligned} x &= 1^{k_d} 2 1^{k_{d-1}} 2 \dots 1^{k_1} 2 1^{k_0}, \\ y &= 2^{l_0} 1 2^{l_1} 1 \dots 2^{l_{d-1}} 1 2^{l_d} \end{aligned}$$

where  $k_0, l_0 \in \mathbb{N}$  and all the other  $k_i$  and  $l_j$  are positive. Thus we have  $\beta(w) = (x, y')$  where

$$y' = 1^{l_d} 2^{l_{d-1}} 2 \dots 1^{l_1} 2^{l_0}.$$

On the other side of the desired equality, we first use Proposition 2.2 to compute

$$\begin{aligned} w &\xrightarrow{\phi^{-1}} 1^{k_0} 2^{l_0+1} 1^{k_1+1} 2^{l_1+1} \dots 2^{l_{d-1}+1} 1^{k_d+1} 2^{l_d} \\ &\xrightarrow{\omega \times \tau} ((k_0, k_1 + 1, k_2 + 1, \dots, k_d + 1), (l_0 + 1, l_1 + 1, \dots, l_{d-1} + 1, l_d)) \\ &\xrightarrow{\sigma \times t} (1^{k_d} 2^{l_{d-1}} 2 \dots 1^{k_1} 2^{l_0}, 1^{l_d} 2^{l_{d-1}} 2 \dots 1^{l_1} 2^{l_0}). \end{aligned}$$

This agrees exactly with the image of  $\beta$  obtained in the previous paragraph.  $\square$

Note that (10) follows as a corollary to part (ii) of this theorem. It is also possible to give  $q$ -analogues of this equality. Let

$$C_{n,d}(q) = \sum_{w \in B_{n-d,d}} q^{\text{inv } w}$$

with  $C_n(q) = C_{2n,n}(q)$ . These  $q$ -Catalan numbers were first defined by Carlitz and Riordan [16]. They have since become the objects of intense study [11, 19, 20, 21, 22, 23, 24, 36, 42, 45, 46, 51, 52, 53]. This is in part because of their connection with the space of diagonal harmonics; see the text of Haglund [41] for more information and references.

We also let

$$c_{n,d}(q, t) = \sum_{w \in B_{n-d,d}} q^{\text{maj } w} t^{\text{des } w}.$$

Comparing with (7) gives  $c_n(q, t) = c_{2n,n}(q, t)$ . We will often use upper and lower case letters for the generating functions of  $\text{inv}$  and  $\text{maj}$ , respectively, over the same set. It will also be convenient to have the notation

$$\delta_{n,d}(q, t) = c_{n,d}(q, t) - c_{n-1,d-1}(q, t).$$

Before stating the  $q$ -analogues, it will be useful to have a result about how our statistics change under the prime operation.

**Lemma 3.4.** *Suppose  $|y| = n$ .*

- (i)  $\text{inv } y' = \text{inv } y$ .
- (ii)  $\text{des } y' = \text{des } y$ .
- (iii)  $\text{maj } y' = n \text{des } y - \text{maj } y$ .

**Proof.** Let  $y = a_1 a_2 \dots a_n$ . Note that  $(i, j)$  is an inversion of  $y$  if and only if  $i < j$  and  $a'_i < a'_j$ , which is if and only if  $(n - j + 1, n - i + 1)$  is an inversion of  $y'$ . All three parts of the lemma now follow.  $\square$

**Proposition 3.5.** *We have*

$$C_n(q) = \sum_{d \geq 0} q^{d^2} C_{n,d}(q)^2,$$

and

$$\begin{aligned} c_n(q, t) = \sum_{d \geq 0} [ & q^n t c_{n-1,d-1}(q, t) c_{n-1,d-1}(q^{-1}, q^{2n}t) + c_{n-1,d-1}(q, t) \delta_{n,d}(q^{-1}, q^{2n}t) \\ & + \delta_{n,d}(q, t) c_{n-1,d-1}(q^{-1}, q^{2n}t) + \delta_{n,d}(q, t) \delta_{n,d}(q^{-1}, q^{2n}t) ]. \end{aligned}$$

**Proof.** Suppose  $w = b_1 b_2 \dots b_{2n} \in B_n$  and let  $x = b_1 \dots b_n$ ,  $y = b_{n+1} \dots b_{2n}$  so that  $\beta(w) = (x, y')$ . If  $x \in B_{n-d,d}$  then  $y \in B_{d,n-d}$  and, by Lemma 3.4(i),

$$\text{inv } w = \text{inv } x + \text{inv } y + d^2 = \text{inv } x + \text{inv } y' + d^2$$

where the  $d^2$  comes from the inversions between twos in  $x$  and ones in  $y$ . Using the fact that  $\beta$  is a bijection, the displayed equation above translates directly into the first desired formula.

To obtain the second formula, it will be convenient to have the characteristic function  $\chi$  which is 1 on true statements and 0 on false ones. Keeping the notation from the first paragraph, we have

$$\begin{aligned} \text{des } w &= \text{des } x + \text{des } y + \chi(b_n > b_{n+1}), \\ \text{maj } w &= \text{maj } x + \text{maj } y + n \text{des } y + n \chi(b_n > b_{n+1}) \end{aligned}$$

where the  $n \text{des } y$  term comes from the fact that  $i$  is a descent of  $y$  if and only if  $n + i$  is a descent of  $w = xy$ . Note also that  $b_n > b_{n+1}$  is equivalent to  $b_n = 2$  and  $b_{n+1} = 1$ . Using Lemma 3.4(ii) and (iii), we obtain

$$\begin{aligned} \text{des } w &= \text{des } x + \text{des } y' + \chi(b_n = b'_{n+1} = 2), \\ \text{maj } w &= \text{maj } x - \text{maj } y' + 2n \text{des } y' + n \chi(b_n = b'_{n+1} = 2). \end{aligned}$$

The proof now breaks down into four cases depending on the values of  $b_n$  and  $b_{n+1}$ . These correspond to the four terms in the expression for  $c_n(q, t)$ . Since the proofs are similar, we will only derive the second of them which corresponds to the

case  $b_n = 2$  and  $b'_{n+1} = 1$ . Using the last pair of displayed equations, we obtain in this case

$$\sum_w q^{\text{maj } w} t^{\text{des } w} = \sum_x q^{\text{maj } x} t^{\text{des } x} \sum_{y'} q^{-\text{maj } y'} (q^{2n} t)^{\text{des } y'}.$$

Now  $x$  is ranging over all elements of  $B(n-d, d)$  which end in a two. But this two does not contribute to maj or des. So removing the two, we get the same sum over all elements of  $B(n-d, d-1)$ . Thus this sum can be replaced by  $c_{n-1, d-1}(q, t)$ . The  $y'$  sum ranges over elements of  $B(n-d, d)$  which end in a 1. Rewriting this as the sum over all of  $B(n-d, d)$  minus the sum over those elements ending in a 2, and then using arguments similar to those concerning the  $x$  sum, gives us the  $\delta_{n,d}(q^{-1}, q^{2n}t)$  factor.  $\square$

Several observations about the expression for  $c_n(q, t)$  are in order. Note that, using the definition of  $\delta_{n,d}(q, t)$ , one can simplify the sum by combining either the 2nd and 4th or the 3rd and 4th terms. But we have chosen the more symmetric form. Note also that such an expression can not be reduced to one in  $q$  alone because of the appearances of  $q^{2n}$  in the second variable. On the other hand, one has the following  $q$ -analogue of (6),

$$c_n(q, 1) = \frac{1}{[n+1]} \begin{bmatrix} 2n \\ n \end{bmatrix},$$

where we have the usual conventions  $[n] = 1 + q + \dots + q^{n-1}$ ,  $[n]! = [1][2] \dots [n]$ , and  $\begin{bmatrix} n \\ k \end{bmatrix} = [n]! / ([k]![n-k]!)$ . No closed form expression is known for  $c_n(q, t)$ .

### 3.3 Extensions

There are analogues of Theorems 3.1 and 3.2 for ballot sequences with any number of ones and twos. Let

$$R_{k,l} = \{\lambda : r_i(\lambda) < 0 \text{ for } 1 \leq i \leq d(\lambda) \text{ and } \lambda \subseteq k \times l\}.$$

We will just state the next result as its proof is similar to the case when  $k = l$ .

**Theorem 3.6.** *Suppose  $k \geq l$ . We have*

$$\phi(B_{k,l}) = \{w \in \Pi(1^k 2^l) : \lambda(w) \in R_{k,l}\}.$$

Also, for  $v$  having the form (4), we have  $\phi(v) \in B_n$  if and only if

(i) for all  $i$  with  $1 \leq i \leq d$

$$\begin{aligned} m_d + m_{d-1} + \dots + m_{d-i+1} &\geq 2i, \\ n_d + n_{d-1} + \dots + n_{d-i+1} + (k-l) &\geq 2i-1, \end{aligned}$$

(ii) as well as

$$\sum_{i=0}^d m_i = k, \quad \text{and} \quad \sum_{i=0}^d n_i = l. \quad \square$$

There is another way to generalize Theorem 3.1. Given  $w \in \{1, 2\}^*$  we let  $e(w)$  be the maximum excess of the number of twos over the number of ones in any prefix of  $w$ . So, if  $w$  has the form (4), then let

$$e_i(v) = n_0 + n_1 + \cdots + n_i - m_0 - m_1 - \cdots - m_i$$

and we have

$$e(w) = \max_{0 \leq i \leq d} e_i.$$

Note that  $w$  is a ballot sequence if and only if  $e(w) \leq 0$ . There is another combinatorial interpretation of  $e(w)$  as follows. One can pair up certain ones and twos (thinking of them as left and right parentheses, respectively) of  $w$  in the usual manner: If a one is immediately followed by a two then they are considered paired. Remove all such pairs from  $w$  and iteratively pair elements of the remaining word. Let  $p(w)$  be the number of pairs in  $w$ . If  $w \in \Pi(1^n 2^n)$ , then it is easy to see by induction on  $n$  that

$$e(w) = n - p(w).$$

We will return to pairings in Section 5. The statistic on integer partitions  $\lambda$  corresponding to  $e(w)$  is the maximum rank

$$r(\lambda) = \max_{1 \leq i \leq d(\lambda)} r_i(\lambda).$$

**Lemma 3.7.** *Suppose  $v \in \{1, 2\}^*$ ,  $\lambda = \lambda(\phi(v))$ , and  $d = d(\lambda)$ . It follows that*

$$e(v) \geq r(\lambda) + 1.$$

*If  $e_d(v) \leq e(v)$  (in particular, if  $v \in \Pi(1^n 2^n)$  is not a ballot sequence) then we have equality.*

**Proof.** As usual, let  $v$  have the form (4). The same reasoning as in the proof of Theorem 3.1 shows that

$$e_i(v) = r_{d-i}(\lambda) + 1 \tag{12}$$

where  $d = \text{des } v = d(\lambda)$  and  $0 \leq i < d$ . It follows that

$$r(\lambda) + 1 = \max_{0 \leq i < d} e_i(v) \leq e(v)$$

giving the desired inequality. It should also be clear why  $e_d(v) \leq e(v)$  implies equality. Finally, suppose  $v \in \Pi(1^n 2^n)$  is not a ballot sequence. It follows that for some  $i > d$  we have  $e_i(v) > 0 = e_d(v)$  and we are done.  $\square$

To state the analogue of Theorem 3.1, let

$$E_{n,k} = \{w \in \Pi(1^n 2^n) : e(w) = k\}$$

and

$$P_{n,k} = \{w \in \Pi(1^n 2^n) : r(\lambda(w)) = k\}.$$

As an immediate corollary of the previous lemma we have the following.

**Theorem 3.8.** *If  $k > 0$  then the pair  $(E_{n,k}, P_{n,k-1})$  is Mahonian.*  $\square$

## 4 Words counted by Fibonacci numbers

We will explore Mahonian pairs constructed from three different sets of sequences enumerated by the Fibonacci numbers. Let the Fibonacci numbers themselves be defined by  $F_0 = F_1 = 1$  and  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 2$ .

The first set of sequences is given by

$$\mathcal{F}_n = \{w \in \{1, 2\}^* : |w| = n \text{ and } w \text{ has no consecutive ones}\}.$$

It is well known and easy to show that  $\#\mathcal{F}_n = F_{n+1}$ . Let

$$f_n(q, t) = \sum_{w \in \mathcal{F}_n} q^{\text{maj } w} t^{\text{des } w}.$$

As with the  $F_n$ , we have a recursion

$$f_n(q, t) = f_{n-1}(q, t) + q^{n-1} t f_{n-2}(q, t)$$

since any  $w \in \mathcal{F}_n$  can be obtained by appending 2 to an element of  $\mathcal{F}_{n-1}$  (which does not change maj or des) or by appending 21 to an element of  $\mathcal{F}_{n-2}$  (which increases maj by  $n-1$  and des by 1). Polynomials satisfying this same recursion were introduced by Carlitz [14, 15] and since studied by a number of authors [26, 25, 27, 28, 39, 38, 56, 57]. However, the initial conditions used for these polynomials is different from ours, and so a different sequence of polynomials is generated. No literature seems to exist about the other  $q$ -analogues of  $F_n$  given by taking the distribution of inv or maj over the various sets considered in this section. It would be interesting to study their properties.

To state our result about what  $\phi$  does to  $\mathcal{F}_n$ , it will be convenient to consider

$$\mathcal{F}_{n,k} = \{w \in \mathcal{F}_n : w \text{ has } k \text{ ones}\}.$$

We also note that if partition  $\lambda$  has Durfee square  $D(\lambda)$  then  $\lambda = D(\lambda) \uplus R(\lambda) \uplus B(\lambda)$  where  $R(\lambda)$  and  $B(\lambda)$  are the connected components of the skew partition  $\lambda/D(\lambda)$  to the right and below  $\lambda$ , respectively. Finally, we let  $\begin{bmatrix} n \\ k \end{bmatrix} = 0$  if  $k < 0$  or  $k > n$ .

**Theorem 4.1.** *We have*

$$\phi(\mathcal{F}_{n,k}) = \{w : \lambda(w) = (\lambda_1, \dots, \lambda_k) \text{ with } \lambda_1 \leq n - k \text{ and } \lambda_k \geq k - 1\}.$$

*It follows that*

$$f_n(q, t) = \sum_{k \geq 0} q^{k(k-1)} t^{k-1} \left( \begin{bmatrix} n - k \\ k - 1 \end{bmatrix} + q^k t \begin{bmatrix} n - k \\ k \end{bmatrix} \right). \quad (13)$$

**Proof.** Suppose  $v \in \mathcal{F}_{n,k}$  and let  $\lambda(\phi(v)) = \lambda$ . Since  $v$  has  $k$  ones and  $n - k$  twos,  $\lambda$  will fit in a  $k \times (n - k)$  rectangle and so  $\lambda_1 \leq n - k$ . Suppose  $v$  has the form (4). So  $v$  having no consecutive ones implies  $m_i = 1$  for  $1 \leq i \leq d = \text{des } v$ . It follows from Proposition 2.2 that

$$\lambda_k = (\# \text{ of initial twos in } \phi(v)) \geq d \geq k - 1$$

where the last inequality follows from the fact that, since the ones in  $v$  are not consecutive, each of them except possibly the last creates a descent. This reasoning is reversible and thus we have our characterization of  $\phi(\mathcal{F}_{n,k})$ .

To get (13), first use Corollary 2.4 to write  $f_n(q, t) = \sum_{\lambda} q^{|\lambda|} t^{d(\lambda)}$  where the sum is over all  $\lambda$  in the description of  $\phi(\mathcal{F}_{n,k})$  for each  $k$ . Since  $\lambda$  has  $k$  parts and  $\lambda_k \geq k - 1$ ,  $d(\lambda) = k - 1$  or  $d(\lambda) = k$ , corresponding to the two terms in the summation. If  $d(\lambda) = k - 1$  then we will have a factor of  $q^{(k-1)^2} t^{k-1}$ . Also in this case  $B(\lambda) = (k - 1)$  and  $R(\lambda) \subseteq (k - 1) \times (n - 2k + 1)$  because  $\lambda \subseteq k \times (n - k)$ . This gives a second factor of  $q^{k-1} \begin{bmatrix} n - k \\ k - 1 \end{bmatrix}$ , and multiplying the two factors gives the first term in (13). The second term is obtained similarly when  $d(\lambda) = k$ .  $\square$

Note that letting  $t = 1$  in (13) and applying one of the usual recursions for the  $q$ -binomial coefficients yields

$$f_n(q, 1) = \sum_{k \geq 0} q^{k(k-1)} \begin{bmatrix} n - k + 1 \\ k \end{bmatrix}.$$

Now letting  $n \rightarrow \infty$  gives, on the right hand side,  $\sum_{k \geq 0} q^{k^2 - k} / (q)_k$  where we are using the Pochhammer symbol  $(q)_k = (1 - q)(1 - q^2) \cdots (1 - q^k)$ . This series was



first studied by Carlitz [12, 13] who related it to the Rogers-Ramanujan identities. Garrett, Ismail, and Stanton [35] generalized the Rogers-Ramanujan identities to sums of the form  $\sum_{k \geq 0} q^{k^2 + mk} / (q)_k$  for any integer  $m$  (so  $m = 0, 1$  are the original identities and  $m = -1$  is the case Carlitz considered). This work has been continued in [43, 44, 5].

To describe  $\phi^{-1}(\mathcal{F}_n)$  we will need a few definitions. A *run* in a word  $w = b_1 b_2 \dots b_n$  is a maximal factor  $r = b_i b_{i+1} \dots b_j$  such that  $b_i = b_{i+1} = \dots = b_j$ . We call  $r$  a *k-run* if the common value of its elements is  $k$ . Furthermore,  $r$  is called the *prefix* or *suffix* run if  $i = 1$  or  $j = n$ , respectively. For example,  $w = 1112212222$  has four runs, the prefix run 111, the suffix run 2222, the 1-run consisting of the rightmost 1, and the 2-run 22.

**Theorem 4.2.** *The set  $\phi^{-1}(\mathcal{F}_n)$  consists of all  $v \in \{1, 2\}^*$  of length  $n$  satisfying the following two conditions.*

- (i) *If  $v$  has a 1-run  $r$  as a prefix or suffix then  $|r| \leq 1$  or  $|r| \leq 2$ , respectively.*
- (ii) *For any run of  $v$  which is neither the prefix nor the suffix, we have  $|r| \leq 2$  for 1-runs and  $|r| \geq 2$  for 2-runs.*

**Proof.** Assume  $v \in \phi^{-1}(\mathcal{F}_n)$  has form (4). By Proposition 2.2,  $\phi(v)$  has no consecutive ones if and only if the following three conditions hold

1.  $m_i - 1 \leq 1$  for  $1 \leq i \leq d$ ,
2.  $n_j - 1 \geq 1$  for  $1 \leq j < d$ , and
3.  $m_0 \leq 1$  with  $n_0 - 1 \geq 1$  if  $m_0 = 1$ .

It is easy to see that these are equivalent to the run conditions. □

Our second set of sequences is closely related to the first. Let

$$\mathcal{G}_n = \{w \in \{1, 2\}^* : |w| = n \text{ and } w \text{ has no consecutive twos}\}.$$

and define generating functions  $g_n(q, t)$  and  $G_n(q)$  analogous to those for  $\mathcal{F}_n$ . Note that by Lemma 3.4(i), the map  $w \mapsto w'$  restricts to an inv-preserving bijection between  $\mathcal{F}_n$  and  $\mathcal{G}_n$ . So  $F_n(q) = G_n(q)$ . There does not seem to be a simple relationship between  $f_n(q, t)$  and  $g_n(q, t)$ .

One can derive analogues for  $\mathcal{G}_n$  of the previous two theorems by modifying the proofs. But we prefer a more elegant approach. Define the *complement* of  $w = b_1 b_2 \dots b_n \in \{1, 2\}^*$  to be

$$c(w) = b'_1 b'_2 \dots b'_n.$$

So  $w'$  is just  $c(w)$  reversed. Also, define the *left rotation* of  $w$  to be

$$\ell(w) = b_2 b_3 \dots b_n b_1.$$

**Lemma 4.3.** *For any  $v \in \{1, 2\}^*$  we have*

$$\phi(c(v)) = \ell(\phi(v)').$$

**Proof.** Let  $v$  have form (4). The proof now breaks up into four cases depending on whether  $m_0$  and  $n_d$  are zero or not. But they are all similar so we will just provide details for  $m_0, n_d > 0$ . On the left-hand side of the equation we have, using Lemma 2.2,

$$\begin{aligned} \phi(c(v)) &= \phi(2^{m_0} 1^{n_0} 2^{m_1} 1^{n_1} \dots 2^{m_d} 1^{n_d}) \\ &= 1^{n_d-1} 2 1^{n_{d-1}-1} 2 \dots 1^{n_0-1} 2 2^{m_0-1} 1 2^{m_1-1} 1 \dots 2^{m_d-1} 1. \end{aligned}$$

Using the same lemma on the right gives

$$\begin{aligned} \ell(\phi(v)') &= \ell(1^{n_d} 2 1^{n_{d-1}-1} 2 \dots 1^{n_0-1} 2^{m_0} 1 2^{m_1-1} 1 2^{m_2-1} \dots 1 2^{m_d-1}) \\ &= 1^{n_d-1} 2 1^{n_{d-1}-1} 2 \dots 1^{n_0-1} 2^{m_0} 1 2^{m_1-1} 1 \dots 2^{m_d-1} 1. \end{aligned}$$

which is clearly equal to what we had before. □

We will use the notation

$$\mathcal{G}_{n,l} = \{w \in \mathcal{G}_n : w \text{ has } l \text{ twos}\}.$$

The proofs of the next two theorems are easily obtained by applying the previous lemma to the results of Theorems 4.1 and 4.2, so we leave them to the reader.

**Theorem 4.4.** *We have*

$$\phi(\mathcal{G}_{n,l}) = \{w \in \Pi(1^{n-l} 2^l) : d(\lambda) \geq l-1, \text{ and } R(\lambda) = \emptyset \text{ where } \lambda = \lambda(w)\}$$

*It follows that*

$$g_n(q, t) = \sum_{l \geq 0} q^{(l-1)^2} t^{l-1} \left( \left[ \begin{matrix} n-l \\ l-1 \end{matrix} \right] + q^{2l-1} t \left[ \begin{matrix} n-l \\ l \end{matrix} \right] \right). \quad \square$$

**Theorem 4.5.** *The set  $\phi^{-1}(\mathcal{G}_n)$  consists of all  $v \in \{1, 2\}^*$  of length  $n$  satisfying the following two conditions.*

- (i) *If  $v$  has a 2-run  $r$  as a prefix or suffix then  $|r| \leq 1$ .*

(ii) For any run of  $v$  which is neither the prefix nor the suffix, we have  $|r| \geq 2$  for 1-runs and  $|r| \leq 2$  for 2-runs.  $\square$

The third set of sequences is

$$\mathcal{H}_n = \{w = b_1 b_2 \dots \in \{1, 2\}^* : \sum_i b_i = n\}.$$

It is easy to prove that  $\#\mathcal{H}_n = F_n$  and this interpretation of the Fibonacci numbers can be made geometric by using tilings. See the book of Benjamin and Quinn [9, p. 1] for examples. Since  $\phi$  preserves the number of ones and twos, it is clear that  $\phi(\mathcal{H}_n) = \mathcal{H}_n$ . We have proved the following result.

**Theorem 4.6.** *The pair  $(\mathcal{H}_n, \mathcal{H}_n)$  is Mahonian. In other words*

$$\sum_{w \in \mathcal{H}_n} q^{\text{maj } w} = \sum_{w \in \mathcal{H}_n} q^{\text{inv } w}. \quad \square$$

## 5 Infinite Mahonian pairs

It is an easy matter to generalize the definition of a Mahonian pair to sets of any cardinality. Call pair  $(S, T)$  with  $S, T \subseteq \mathbb{P}^*$  *Mahonian* if there is a bijection  $\alpha : S \rightarrow T$  such that  $\text{maj } v = \text{inv } \alpha(v)$  for all  $v \in S$ . In this section we will study infinite Mahonian pairs connected to ballot sequences. In particular, we will be able to show that a bijection of Corteel-Savage-Venkatraman [29] between partitions with all ranks positive and partitions with no part equal to 1 is essentially conjugation by  $\phi$  of the map used by Greene-Kleitman to obtain a symmetric chain decomposition of the Boolean algebra [40].

Our starting point is an observation made independently by Andrews [3] and Erdős-Richmond [32]. Let  $P$  be the set of all integer partitions. Define

$$R_{\geq t} = \{\lambda \in P : r_i(\lambda) \geq t \text{ for all } i\}$$

and similarly for  $R_{\leq t}$ . Also, let

$$P_{\neq t} = \{\lambda \in P : \lambda_i \neq t \text{ for all } i\}.$$

**Theorem 5.1** ([3, 32]). *We have*

$$\sum_{\lambda \in R_{\geq 1}} q^{|\lambda|} = \sum_{\lambda \in P_{\neq 1}} q^{|\lambda|}. \quad \square$$

We wish to make a couple of comments about this theorem. First of all, it is well known that

$$\sum_{\lambda \in P_{\neq 1}} q^{|\lambda|} = \prod_{i \geq 2} \frac{1}{1 - q^i}. \quad (14)$$

Secondly, Theorem 5.1 is a special case of an earlier result of Andrews [2] as generalized by Bressoud [10]. (Andrews had the restriction that  $M$  below must be odd.)

**Theorem 5.2** ([2, 10]). *Let  $M, r$  be integers satisfying  $0 < r < M/2$ . The number of partitions of  $n$  whose ranks lie in the interval  $[-r + 2, M - r - 2]$  equals the number of partitions of  $n$  with no part congruent to  $0, r$ , or  $-r$  modulo  $M$ .  $\square$*

Note that if  $M = n + 2$  and  $r = 1$  then one recovers Theorem 5.1. And for  $M = 5$  and  $r = 1, 2$  one obtains the Rogers-Ramanujan identities.

In order to involve Foata's map, we will need to associate with each partition  $\lambda$  the word  $w(\lambda)$  gotten by recording the north and east steps along the southeast boundary of  $\lambda$ 's Ferrers diagram with ones and twos, respectively. To illustrate,  $w(3, 2, 2) = 221121$  as can be seen from Figure 1. Clearly  $w = w(\lambda)$  for some nonempty  $\lambda$  if and only if  $w \in 2\{1, 2\}^*1$ , i.e.,  $w$  begins with a two and ends with a one. We will often abuse notation and write things like  $\phi(\lambda)$  for the more cumbersome  $\phi(w(\lambda))$ . To state our first result about infinite Mahonian pairs, we will use the notation

$$W_w = \{1, 2\}^* w \uplus \{\epsilon\}$$

for any word  $w$ , and

$$B_w = \{w \in W_w : w \text{ is a ballot sequence}\}.$$

Furthermore, for any set  $\Lambda$  of partitions we let  $\Lambda' = \{\lambda' : \lambda \in \Lambda\}$ .

**Theorem 5.3.** *We have*

$$\phi^{-1}(P) = W_{21}, \quad \phi^{-1}(R'_{\geq 1}) = B_{21}, \quad \text{and} \quad \phi^{-1}(P'_{\neq 1}) = W_{121}.$$

*It follows that  $(S, T) = (W_{21}, P)$ ,  $(B_{21}, R'_{\geq 1})$ , and  $(W_{121}, P'_{\neq 1})$  are Mahonian pairs. Furthermore, for any of these pairs  $(S, T)$  we have*

$$\sum_{w \in S} q^{\text{maj } w} t^{\text{des } w} z^{e(w)} = \sum_{\lambda \in T} q^{|\lambda|} t^{d(\lambda)} z^{r(\lambda)}. \quad (15)$$

*We also have*

$$\sum_{w \in B_{21}} q^{\text{maj } w} = \sum_{w \in W_{121}} q^{\text{maj } w} = \prod_{i \geq 2} \frac{1}{1 - q^i}.$$

**Proof.** The fact that  $\phi^{-1}(P) = W_{21}$  follows from Lemma 2.3 and the observation above that  $w = w(\lambda)$  precisely when  $w \in 2\{1, 2\}^*1$ . Note that if  $w \in W_{21}$  then  $e_d(w) < e_{d-1}(w)$ , and this implies equality in Lemma 3.7. Combining this with Corollary 2.4 yields (15) for the case  $(W_{21}, P)$ . Since the other two pairs are formed from subsets of  $W_{21}$  and  $P$ , the same equation will hold for them once we show that they are mapped in the desired way by  $\phi$ .

To prove that  $\phi^{-1}(R'_{\geq 1}) = B_{21}$ , note that  $R'_{\geq 1} = R_{\leq -1}$ . Let us demonstrate that  $\phi(B_{21}) \subseteq R_{\leq -1}$ . Suppose that  $v \in B_{21} - \{\epsilon\}$  is of the form (4) and let  $w = \phi(v)$ . Then by Lemma 2.1(iii) we have  $\phi(v) \in 2\{1, 2\}^*1$  and so there is a partition  $\lambda$  with  $w(\lambda) = w$ . Since  $v$  is a ballot sequence we have  $e(v) \leq 0$ . It follows from Lemma 3.7 that  $r(\lambda) \leq -1$  giving the desired set containment.

For the reverse containment, start with a nonempty partition  $\lambda \in R_{\leq -1}$  and let  $w = w(\lambda)$ . From the first paragraph of the proof we already know  $v = \phi^{-1}(\lambda) \in W_{21}$ . Using (12), we see that  $\lambda \in R_{\leq -1}$  forces  $e_i(v) \leq 0$  for  $i < d = \text{des } v$ . And since  $v = u21$  for some  $u$ , we have that  $e_d(v) = e_{d-1}(v) - 1$ . It follows that  $e(v) \leq 0$  which is equivalent to  $v$  being a ballot sequence. Thus  $v \in B_{21}$  which is what we needed to prove.

The demonstration of  $\phi^{-1}(P'_{\neq 1}) = W_{121}$  is similar to the one just given, using the fact that  $\lambda \in P'_{\neq 1}$  if and only if  $w = w(\lambda) \in 2\{1, 2\}^*11$ . The final statement about generating functions now follows from Lemma 3.4(i), Theorem 5.1, and equation (14).  $\square$

Corteel, Savage, and Venkatraman [29] gave a bijective proof of Theorem 5.1. Using  $\phi$ , we can relate their function to the Greene-Kleitman symmetric chain decomposition of a Boolean algebra [40]. For simplicity in comparing with our results thus far, we will describe the conjugate of their map. For a partition  $\lambda$  it will be convenient to define  $\delta(\lambda) = \lambda_1 - \lambda_2$  and let

$$D_t = \{\lambda \in P : \delta(\lambda) = t\}.$$

Note that  $P'_{\neq 1} = D_0$ . Define a map  $CSV : D_0 \rightarrow R_{\leq -1}$  by the following algorithm:


CSV1 Input  $\lambda \in D_0$ .

CSV2 While  $r(\lambda) \geq 0$  do

- (a) Let  $i$  be the maximum index such that  $r_i(\lambda) = r(\lambda)$ .
- (b) Remove a part of size  $i$  from  $\lambda'$ , add a part of size  $i-1$  to  $\lambda$ , and increase the size of  $\lambda_1$  by one.

CSV3 Output  $\lambda$ .

We have written out the complete algorithm for computing  $CSV(8, 8, 6, 5, 2, 1)$  below where dots rather than boxes have been used for the Ferrers diagrams. At each stage, the rank vector  $\rho(\lambda) = [r_1(\lambda), \dots, r_d(\lambda)]$  is displayed along with the maximum  $r = r(\lambda)$  and the largest index  $i = i(\lambda)$  where  $r$  is achieved. For future reference, we have also displayed  $w = w(\lambda)$ ,  $v = \phi^{-1}(w)$ , and  $\varepsilon(v) = [e_0(v), \dots, e_d(v)]$ .



Let  $\kappa$  denote the map obtained by going once through the loop at CSV2. Now for an arbitrary partition  $\lambda$ ,  $\kappa(\lambda)$  may not be well defined since it may not be possible to find a part of size  $i(\lambda)$  in  $\lambda'$ . But it is shown in [29] that if one starts with  $\lambda \in D_0$  then such a part must exist while  $r(\lambda) \geq 0$ . One also needs to worry about termination of the algorithm. But in [29] it is shown that while  $r(\lambda) \geq 0$  we

have  $r(\kappa(\lambda)) \leq r(\lambda) - 1$  with equality if  $r(\lambda) > 0$ . So

$$CSV(\lambda) = \kappa^{r+1}(\lambda) \tag{16}$$

where  $r = r(\lambda)$ .

In their seminal paper [40], Green and Kleitman gave a symmetric chain decomposition of the Boolean algebra  $\mathcal{B}_n$  of all subsets of  $\{1, \dots, n\}$  as follows. Represent an element  $S \in \mathcal{B}_n$  as a word  $w = w_1 \dots w_n \in \{1, 2\}^*$  where  $w_i = 2$  if and only if  $i \in S$ . Pair ones and twos in  $w$  as described near the end of Section 3. Note that the unpaired twos must precede the unpaired ones. Define a map  $\gamma$  on words with at least one unpaired two by

$$\gamma(w) = w \text{ with the rightmost unpaired two changed to a one.}$$

Note that, by the choice of the changed two,  $\gamma(w)$  has the same pairs as  $w$ . The chains in the decomposition are all those of the form  $w, \gamma(w), \gamma^2(w), \dots, \gamma^t(w)$  where the unpaired elements of  $w$  are all twos and  $t$  is the number of such elements (so the unpaired elements of  $\gamma^t(w)$  are all ones).

Define a map  $GK : W_{121} \rightarrow B_{21}$  as follows. If  $v = x121 \in W_{121}$  has  $t$  unpaired twos (note that they must all be in  $x$ ), then let

$$GK(v) = \gamma^t(x)12^{t+1}1. \tag{17}$$

**Theorem 5.4.** *The map  $GK : W_{121} \rightarrow B_{21}$  is well defined and bijective.*

**Proof.** For  $GK$  to be well defined we must check  $GK(v) \in B_{21}$  for  $v \in W_{121}$ . It is clear from (17) that  $v \in \{1, 2\}^*21$ . Also,  $GK(v)$  has no unmatched twos since  $\gamma^t(x)$  has replaced the  $t$  unmatched twos in  $x$  with ones, and so they can be used (along with the penultimate one) to match the twos in  $2^{t+1}$ . But having no unmatched twos is equivalent to being a ballot sequence and so  $GK(v) \in B_{21}$ .

To prove bijectivity, we construct an inverse map. Given  $w \in B_{21}$  we write  $w = y12^{t+1}1$  for some  $t \geq 0$ . Since  $w$  is a ballot sequence,  $t$  of the twos in the last 2-run must be matched with ones in  $y$  which are unmatched in  $y$  itself. So  $x = \gamma^{-t}(y)$  is well defined and we can set  $v = x121$ . It is straightforward to check that  $GK(v) = w$  and that this construction provides an inverse to  $GK$ .  $\square$

We now have all the tools in hand to demonstrate the relationship between  $CSV$  and  $GK$ .

**Theorem 5.5.** *We have*

$$CSV = \phi \circ GK \circ \phi^{-1}. \tag{18}$$

**Proof.** Consider  $\lambda \in D_0$  and  $v = \phi^{-1}(\lambda)$ . If  $\lambda \in R_{\leq -1}$  as well then  $CSV(\lambda) = \lambda$ . We also have  $v \in W_{121} \cap B_{21}$ . It follows that  $v = x121$  where  $x$  is a ballot sequence and so has no unpaired twos. This implies that  $GK(v) = v$  and both sides of (18) agree, as desired.

From now on we can assume  $\lambda \notin R_{\leq -1}$  and thus  $r = r(\lambda) \geq 0$ . Appealing to Lemma 3.7, we have  $e(v) \geq 1$ . In this case we will show that if  $t$  is the number of unpaired ones in  $v$  then  $r + 1 = t$ . Note that for any  $v$ , if  $e_i(v)$  is a positive left-right maximum in the sequence  $e_1(v), e_2(v), \dots$  then the  $i$ th 2-run in  $v$  has exactly  $e_i(v) - e_j(v)$  unpaired twos, where  $e_j(v)$  is the positive left-right maximum just before  $e_i(v)$  (or 0 if there is no such previous value). Since  $e(v) > 0$  we know that a positive left-right maximum exists and thus  $e(v) = t$ , the number of unpaired twos. We also have that  $v \in W_{121}$  implying that  $e_d(v) = e_{d-1}(v) - 1$  and so  $e_d(v) < e(v)$ . Applying Lemma 3.7 again shows that  $r + 1 = e(v) = t$  as claimed.

We now consider a variant  $\bar{\gamma}$  of  $\gamma$  as follows. Let  $u = x12^m1$  where  $x$  has at least one unpaired two and  $m \geq 1$ . Define

$$\bar{\gamma}(u) = \gamma(x)12^{m+1}1.$$

By the definitions (16) and (17), we have  $GK(v) = \bar{\gamma}^t(v)$  and  $CSV(\lambda) = \kappa^{r+1}(\lambda)$  where  $t = r + 1$ . So the theorem will follow if we can show that  $\kappa = \phi \circ \bar{\gamma} \circ \phi^{-1}$  or, equivalently, that  $\kappa(\phi(y)) = \phi(\bar{\gamma}(y))$  whenever  $y = \bar{\gamma}^j(v)$  for some  $j < t$ .

To set notation, let  $\mu = \lambda(\phi(y))$ ,  $z = \bar{\gamma}(y)$ , and  $\nu = \lambda(\phi(z))$ . We can complete the proof by showing that  $\kappa(\mu) = \nu$ . Let  $y$  have the form (4). Since  $y = \bar{\gamma}^j(v)$  where  $v \in W_{121}$  we see that  $n_d = 0$ ,  $m_d = 1$ , and  $n_{d-1} = j + 1$ . We must also locate the rightmost unpaired two in  $y$ . As in the CSV algorithm, let  $i = i(\mu)$  denote the maximum index such that  $r_i(\lambda) = r(\lambda)$ . Using equation (12) and the fact that  $e_d(y) < e(y)$  since  $y$  ends with a single one, we see that  $d - i$  is the minimum index such that  $e_{d-i}(y) = e(y)$ . So, by the description of the location of the unpaired twos in the second paragraph of this proof, the rightmost such must lie in the  $(d - i)$ th run. Combining all the information gathered thus far and using the definition of  $\bar{\gamma}$  permits us to write

$$\begin{aligned} y &= p2^{n_{d-i}}1^{m_{d-i+1}}q12^{j+1}1, \\ z &= p2^{n_{d-i}-1}1^{m_{d-i+1}+1}q12^{j+2}1 \end{aligned}$$

for some words  $p, q$ .

We now wish to apply  $\phi$  using Proposition 2.2. To do so, we must worry about whether  $n_{d-i} - 1 = 0$  in  $z$ . But in that case  $n_{d-i} = 1$  and the only way a run consisting of a single two could have that two unpaired is if it is the initial run. Since this is also the rightmost unpaired two, it must be the only unpaired two



and so  $j = t - 1$ . The proof in this special case is similar to the one for  $j < t - 1$  which we will present, leaving the last value of  $j$  for the reader.

For  $j < t - 1$  we can write

$$\begin{aligned}\phi(y) &= \bar{p}1^{m_{d-i+1}-1}\bar{q}2^{n_{d-i}-1}\bar{s}12^j1, \\ \phi(z) &= \bar{p}1^{m_{d-i+1}}\bar{q}2^{n_{d-i}-2}\bar{s}12^{j+1}1\end{aligned}$$

for certain  $\bar{p}, \bar{q}, \bar{s}$ . So to pass from  $\phi(y)$  to  $\phi(z)$  we have done the following. We have added a one to the run  $1^{m_{d-i+1}-1}$  in  $\phi(y)$ . By Proposition 2.2, there are exactly  $i - 1$  twos before this run and so this corresponds to adding a part of size  $i - 1$  to  $\mu$  when forming  $\nu$ . Similarly, subtracting a two from the run  $2^{n_{d-i}-1}$  corresponds to deleting a part of size  $i$  from  $\mu'$ . And a two is added to the final 2-run, which corresponds to increasing the size of  $\mu_1$  by one. In summary, we have  $\kappa(\mu) = \nu$  and are done.  $\square$

## 6 The past, the future, and open problems

### 6.1 Lucanomialials

We will describe the genesis of the notion of a Mahonian pair. This is not only for historical reasons, but also because the original problem which lead us to this definition is still unsolved.

Let  $s, t$  be variables and define a sequence of polynomials  $\{n\}$  in  $s$  and  $t$  by  $\{0\} = 0$ ,  $\{1\} = 1$ , and

$$\{n\} = s\{n-1\} + t\{n-2\} \tag{19}$$

for  $n \geq 2$ . When  $s, t$  are integers, the arithmetic properties of this sequence were studied by Lucas [47, 49, 48]. Given integers  $0 \leq k \leq n$ , define the corresponding *lucanomial coefficient*

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \frac{\{n\}!}{\{k\}!\{n-k\}!}.$$

Note that one obtains the fibonomial coefficients or  $q$ -binomial coefficients by specializing to  $s = t = 1$  or to  $s = [2]$  and  $t = -q$ , respectively. It is not hard to show that these rational functions are actually polynomials in  $s, t$  with coefficients in  $\mathbb{N}$ . In [55], we were able to give a simple combinatorial interpretation of the lucanomial coefficients using tilings of partitions contained in a  $k \times (n - k)$  rectangle. (Earlier, more complicated, interpretations were given by Gessel-Viennot [37] and Benjamin-Plott [7, 8].)

Upon hearing a talk on this subject at the 2010 Mathfest, Lou Shapiro asked the following natural question. Define an  $s, t$ -analogue of the Catalan numbers by

$$C_{\{n\}} = \frac{1}{\{n+1\}} \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\}.$$

**Question 6.1** (Shapiro). *Is  $C_{\{n\}} \in \mathbb{N}[s, t]$ ? If so, is there a simple combinatorial interpretation?*

Ekhad [31] has pointed out that the answer to the first question is "yes" since one can show that

$$C_{\{n\}} = \left\{ \begin{matrix} 2n \\ n \end{matrix} \right\} + t \left\{ \begin{matrix} 2n \\ n-1 \end{matrix} \right\}.$$

However, a nice combinatorial interpretation is elusive. While trying to find such an interpretation, we looked at statistics on ballot sequences (being one of the most common objects associated with  $C_n$ ) and partitions with all ranks positive sitting inside an  $n \times n$  rectangle (since Savage had done previous work with such partitions). It was noted that  $\text{maj}$  on the former was equidistributed with  $\text{area}$  on the latter, and the concept of a Mahonian pair was born.

## 6.2 Larger alphabets, other maps

It would be very interesting to obtain results about Mahonian pairs using sets outside of  $\{1, 2\}^*$ . For example, one could consider ballot sequences over a larger alphabet. The difficulty here is what the analogue of the Ferrers diagram  $\lambda(w)$  should be.

Another possibility would be to use maps other than Foata's. There is another well-known bijection  $\psi : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$  on the symmetric group which exchanges  $\text{maj}$  and  $\text{inv}$ . The *inv labeling* of  $w = a_1 \dots a_n \in \mathfrak{S}_n$  labels the spaces between the  $a_i$  as  $0, \dots, n$  from right to left. For example,  $w = 31524$  has  $\text{inv labeling } {}_5 3_4 1_3 5_2 2_1 4_0$ . Clearly, if  $x$  is the word obtained by inserting  $n+1$  in the  $i$ th space of the  $\text{inv labeling}$  then  $\text{inv } x = \text{inv } w + i$ . Also define the *maj labeling* by first labeling the last space and the descent spaces  $0, \dots, \text{des } w$  from right to left, and then labeling the first space and the ascent spaces  $\text{des } w + 1, \dots, n$  from left to right. In our example, we would get  ${}_3 3_2 1_4 5_1 2_5 4_0$ . It is not hard to check that if  $y$  is the result of putting  $n+1$  in the  $i$ th  $\text{maj space}$  then  $\text{maj } y = \text{maj } w + i$ .

The map  $\psi$  is now defined as follows. Given  $v \in \mathfrak{S}_n$ , consider the sequence  $1 = v_1, v_2, \dots, v_n = v$  where  $v_i$  is the subpermutation of  $v$  on the numbers  $1, \dots, i$ . Now construct  $1 = w_1, w_2, \dots, w_n = \psi(v)$  inductively. For  $j \geq 2$ , if  $v_j$  is obtained from  $v_{j-1}$  by inserting  $j$  in the  $i$ th  $\text{maj space}$ , then  $w_j$  is obtained from  $w_{j-1}$  by inserting  $j$  in the  $i$ th  $\text{inv space}$ . From the definitions we have  $\text{maj } v = \text{inv } \psi(v)$ .

**Question 6.2.** *Can  $\psi$  be used to construct interesting Mahonian pairs?*

One natural subset of  $\mathfrak{S}_n$  to look at would be the up-down (or alternating) permutations  $A_n$  because of the fact that they have constant maj. However, we were not able to characterize either  $\phi(A_n)$  or  $\psi(A_n)$ .

Another place to look would be at sets of permutations determined by pattern avoidance. If  $\pi \in \mathfrak{S}_k$  then we say  $\sigma \in \mathfrak{S}_n$  contains  $\pi$  as a *pattern* if there is a subpermutation of  $\sigma$  order isomorphic to  $\pi$ . If  $\sigma$  does not contain  $\pi$  as a pattern then we say it *avoids*  $\pi$ . We let  $\text{Av}_n(\pi)$  denote the set of such patterns in  $\mathfrak{S}_n$ . Since  $\#\text{Av}_n(\pi) = C_n$  for any  $\pi \in \mathfrak{S}_3$ , the hope was that there would be some connection with the Mahonian pairs associated with ballot sequences. Unfortunately, looking at the distributions for small  $n$  showed no possible pairs among the  $\text{Av}_n(\pi)$  where  $\pi \in \mathfrak{S}_3$ .

However, another phenomenon manifested itself. There were pairs  $\pi, \sigma$  such that the distribution of  $\text{inv}$  over both  $\text{Av}_n(\pi)$  and  $\text{Av}_n(\sigma)$  were the same, and similarly for  $\text{maj}$ . This lead us to the following refinement of the traditional notion of Wilf equivalence. (We say  $\pi$  and  $\sigma$  are *Wilf equivalent* if  $\#\text{Av}_n(\pi) = \#\text{Av}_n(\sigma)$  for all  $n \geq 0$ .) Let  $\text{st}$  be any statistic on permutations. Call  $\pi$  and  $\sigma$  *st-Wilf equivalent* if  $\text{st}$  is equidistributed over  $\text{Av}_n(\pi)$  and  $\text{Av}_n(\sigma)$ . So  $\text{st-Wilf}$  equivalence implies Wilf-equivalence, but not conversely. We are currently in the process of studying this concept.

### 6.3 Eulerian pairs

There is another famous pair of equidistributed statistics. An *excedance* in a permutation  $w = a_1 \dots a_n$  is an index  $i$  such that  $a_i > i$ . Let  $\text{exc } w$  be the number of excedances. It is well known that  $\text{des}$  and  $\text{exc}$  are equidistributed over  $\mathfrak{S}_n$  and any statistic with this distribution is called *Eulerian*. It is easy to extend the definition of excedance to any  $w \in \mathbb{P}^*$ . Let  $x = b_1 \dots b_n$  be the weakly increasing rearrangement of  $w$ . Then an excedance of  $w$  is an index  $i$  with  $a_i > b_i$ . The next definition should come as no surprise. The subsets  $S, T \subseteq \mathbb{P}^*$  form an *Eulerian pair*,  $(S, T)$ , if there is a bijection  $\alpha : S \rightarrow T$  such that

$$\text{exc } v = \text{des } \alpha(v)$$

for all  $v \in S$ . The study of Eulerian pairs could be every bit as rich as that of their Mahonian cousins and we will be investigating their properties.

*Acknowledgment.* We would like to thank Dominique Foata, Ira Gessel, and Drew Sills for useful discussions and references.

## References

- [1] AISSSEN, M. Variations on a theme of Pólya. In *Second International Conference on Combinatorial Mathematics (New York, 1978)*, vol. 319 of *Ann. New York Acad. Sci.* New York Acad. Sci., New York, 1979, pp. 1–6.
- [2] ANDREWS, G. E. Sieves for theorems of Euler, Ramanujan, and Rogers. In *The Theory of Arithmetic Functions*, vol. 251 of *Lecture Notes in Mathematics*. Springer-Verlag, 1971, pp. 1–20.
- [3] ANDREWS, G. E. On the difference of successive Gaussian polynomials. *J. Statist. Plann. Inference* 34, 1 (1993), 19–22.
- [4] ANDREWS, G. E. *The theory of partitions*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1998. Reprint of the 1976 original.
- [5] ANDREWS, G. E., KNOPFMACHER, A., AND PAULE, P. An infinite family of Engel expansions of Rogers-Ramanujan type. *Adv. in Appl. Math.* 25, 1 (2000), 2–11.
- [6] BARNES, T. M., AND SAVAGE, C. D. A recurrence for counting graphical partitions. *Electron. J. Combin.* 2 (1995), Research Paper 11, approx. 10 pp. (electronic).
- [7] BENJAMIN, A. T., AND PLOTT, S. S. A combinatorial approach to Fibonomial coefficients. *Fibonacci Quart.* 46/47, 1 (2008/09), 7–9.
- [8] BENJAMIN, A. T., AND PLOTT, S. S. A combinatorial approach to Fibonomial coefficients: errata. *Fibonacci Quart.* 48, 3 (2010), 276.
- [9] BENJAMIN, A. T., AND QUINN, J. J. *Proofs that really count*, vol. 27 of *The Dolciani Mathematical Expositions*. Mathematical Association of America, Washington, DC, 2003. The art of combinatorial proof.
- [10] BRESSOUD, D. Extension of the partition sieve. *J. Number Theory* 12 (1980), 87–100.
- [11] BUTLER, L. M., AND FLANIGAN, W. P. A note on log-convexity of  $q$ -Catalan numbers. *Ann. Comb.* 11, 3-4 (2007), 369–373.
- [12] CARLITZ, L. A note on the Rogers-Ramanujan identities. *Math. Nachr.* 17 (1958), 23–26.

- [13] CARLITZ, L. Some formulas related to the Rogers-Ramanujan identities. *Ann. Mat. Pura Appl. (4)* 47 (1959), 243–251.
- [14] CARLITZ, L. Fibonacci notes. III.  $q$ -Fibonacci numbers. *Fibonacci Quart.* 12 (1974), 317–322.
- [15] CARLITZ, L. Fibonacci notes. IV.  $q$ -Fibonacci polynomials. *Fibonacci Quart.* 13 (1975), 97–102.
- [16] CARLITZ, L., AND RIORDAN, J. Two element lattice permutation numbers and their  $q$ -generalization. *Duke Math. J.* 31 (1964), 371–388.
- [17] CHEN, W. Y. C., AND HOU, Q.-H. Factors of the Gaussian coefficients. *Discrete Math.* 306, 13 (2006), 1446–1449.
- [18] CHEN, W. Y. C., WANG, C. J., AND WANG, L. X. W. The limiting distribution of the coefficients of the  $q$ -Catalan numbers. *Proc. Amer. Math. Soc.* 136, 11 (2008), 3759–3767.
- [19] CIGLER, J. Operatormethoden für  $q$ -Identitäten. IV. Eine Klasse von  $q$ -Gould-Polynomen. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 205 (1996), 169–174 (1997).
- [20] CIGLER, J. Operatormethoden für  $q$ -Identitäten. V.  $q$ -Catalan-Bäume. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 205 (1996), 175–182 (1997).
- [21] CIGLER, J. Operatormethoden für  $q$ -Identitäten. VI. Geordnete Wurzelbäume und  $q$ -Catalan-Zahlen. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 206 (1997), 253–266 (1998).
- [22] CIGLER, J. Operatormethoden für  $q$ -Identitäten. VII.  $q$ -Catalan-Determinanten. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 208 (1999), 123–142 (2000).
- [23] CIGLER, J.  $q$ -Catalan- und  $q$ -Motzkinzahlen. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 208 (1999), 3–20 (2000).
- [24] CIGLER, J. Einige  $q$ -Analoge der Catalan-Zahlen. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 209 (2000), 19–46 (2001).
- [25] CIGLER, J. A new class of  $q$ -Fibonacci polynomials. *Electron. J. Combin.* 10 (2003), Research Paper 19, 15 pp. (electronic).

- [26] CIGLER, J.  $q$ -Fibonacci polynomials. *Fibonacci Quart.* 41, 1 (2003), 31–40.
- [27] CIGLER, J. Some algebraic aspects of Morse code sequences. *Discrete Math. Theor. Comput. Sci.* 6, 1 (2003), 55–68 (electronic).
- [28] CIGLER, J.  $q$ -Fibonacci polynomials and the Rogers-Ramanujan identities. *Ann. Comb.* 8, 3 (2004), 269–285.
- [29] CORTEEL, S., SAVAGE, C. D., AND VENKATRAMAN, R. A bijection for partitions with all ranks at least  $t$ . *J. Combin. Theory Ser. A* 83, 2 (1998), 202–220.
- [30] DYSON, F. J. Some guesses in the theory of partitions. *Eureka (Cambridge)* 8 (1944), 10–15.
- [31] EKHAD, S. The Sagan-Savage Lucas-Catalan polynomials have positive coefficients. Preprint <http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/bruce.html>.
- [32] ERDŐS, P., AND RICHMOND, L. B. On graphical partitions. *Combinatorica* 13, 1 (1993), 57–63.
- [33] FOATA, D. On the Netto inversion number of a sequence. *Proc. Amer. Math. Soc.* 19 (1968), 236–240.
- [34] FÜRLINGER, J., AND HOFBAUER, J.  $q$ -Catalan numbers. *J. Combin. Theory Ser. A* 40, 2 (1985), 248–264.
- [35] GARRETT, K., ISMAIL, M. E. H., AND STANTON, D. Variants of the Rogers-Ramanujan identities. *Adv. in Appl. Math.* 23, 3 (1999), 274–299.
- [36] GESSEL, I. A noncommutative generalization and  $q$ -analog of the Lagrange inversion formula. *Trans. Amer. Math. Soc.* 257, 2 (1980), 455–482.
- [37] GESSEL, I., AND VIENNOT, G. Binomial determinants, paths, and hook length formulae. *Adv. in Math.* 58, 3 (1985), 300–321.
- [38] GOYT, A. M., AND MATHISEN, D. Permutation statistics and  $q$ -Fibonacci numbers. *Electron. J. Combin.* 16, 1 (2009), Research Paper 101, 15.
- [39] GOYT, A. M., AND SAGAN, B. E. Set partition statistics and  $q$ -Fibonacci numbers. *European J. Combin.* 30, 1 (2009), 230–245.
- [40] GREENE, C., AND KLEITMAN, D. J. Strong versions of Sperner’s theorem. *J. Combinatorial Theory Ser. A* 20, 1 (1976), 80–88.

- [41] HAGLUND, J. *The  $q,t$ -Catalan numbers and the space of diagonal harmonics*, vol. 41 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2008. With an appendix on the combinatorics of Macdonald polynomials.
- [42] INKERI, K. On Catalan's conjecture. *J. Number Theory* 34, 2 (1990), 142–152.
- [43] ISMAIL, M. E. H., PRODINGER, H., AND STANTON, D. Schur's determinants and partition theorems. *Sém. Lothar. Combin.* 44 (2000), Art. B44a, 10 pp. (electronic).
- [44] ISMAIL, M. E. H., AND STANTON, D. Tribasic integrals and identities of Rogers-Ramanujan type. *Trans. Amer. Math. Soc.* 355, 10 (2003), 4061–4091 (electronic).
- [45] KIM, J. H., AND PITTEL, B. Confirming the Kleitman-Winston conjecture on the largest coefficient in a  $q$ -Catalan number. *J. Combin. Theory Ser. A* 92, 2 (2000), 197–206.
- [46] KRATTENTHALER, C. Counting lattice paths with a linear boundary. II.  $q$ -ballot and  $q$ -Catalan numbers. *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 198, 4-7 (1989), 171–199.
- [47] LUCAS, E. Theorie des Fonctions Numeriques Simplement Periodiques. *Amer. J. Math.* 1, 2 (1878), 184–196.
- [48] LUCAS, E. Theorie des Fonctions Numeriques Simplement Periodiques. *Amer. J. Math.* 1, 4 (1878), 289–321.
- [49] LUCAS, E. Theorie des Fonctions Numeriques Simplement Periodiques. [Continued]. *Amer. J. Math.* 1, 3 (1878), 197–240.
- [50] MACMAHON, P. A. *Collected papers. Vol. I*. MIT Press, Cambridge, Mass., 1978. Combinatorics, Mathematicians of Our Time, Edited and with a preface by George E. Andrews, With an introduction by Gian-Carlo Rota.
- [51] MAZZA, C., AND PIAU, D. Products of correlated symmetric matrices and  $q$ -Catalan numbers. *Probab. Theory Related Fields* 124, 4 (2002), 574–594.
- [52] RANDRIANARIVONY, A. Fractions continues,  $q$ -nombres de Catalan et  $q$ -polynômes de Genocchi. *European J. Combin.* 18, 1 (1997), 75–92.

- [53] RAWLINGS, D. The Euler-Catalan identity. *European J. Combin.* 9, 1 (1988), 53–60.
- [54] ROUSSEAU, C. C., AND ALI, F. A note on graphical partitions. *J. Combin. Theory Ser. B* 64, 2 (1995), 314–318.
- [55] SAGAN, B. E., AND SAVAGE, C. D. Combinatorial interpretations of binomial coefficient analogues related to Lucas sequences. Preprint [arXiv:0911:3159](https://arxiv.org/abs/0911.3159).
- [56] SHATTUCK, M. A., AND WAGNER, C. G. Parity theorems for statistics on domino arrangements. *Electron. J. Combin.* 12 (2005), Note 10, 13 pp. (electronic).
- [57] SHATTUCK, M. A., AND WAGNER, C. G. A new statistic on linear and circular  $r$ -mino arrangements. *Electron. J. Combin.* 13, 1 (2006), Research Paper 42, 15 pp. (electronic).
- [58] STANLEY, R. P. *Enumerative Combinatorics. Vol. 1*, vol. 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.